

LIGHT *Sheer*[™] Laser Hair Removal

Pre- and Post-treatment Patient Instructions

Prior to Treatment

- 1 Do not pluck, wax, use a depilatory or undergo electrolysis in the areas you wish to have treated for 6 weeks prior to laser hair removal.
- 2 Do not tan the areas to be treated for 4 weeks prior to treatment.
- 3 Avoid using self-tanning products for 2 weeks prior to treatment.

After Treatment

- 4 Some redness and swelling in the area is normal after treatment and may feel similar to a sunburn. This should resolve within several hours to several days after treatment.
- 5 Gently clean area twice daily.
- 6 Avoid irritants (glycolics, retinoids etc.) for seven days after treatment.
- 7 Apply sunscreen for 6 weeks over the treated area.

DRUGS THAT CAUSE PHOTOSENSITIVITY

Anticancer Drugs

Dacarbazine
Fluorouracil
Flutamide
Methotrexate
Vinblastine

Antihistamines

Cyproheptadine
Diphenhydramine

Antimicrobials

Ciprofloxacin
Clofazimine
Dapsone
Demeclocycline
Doxycycline
Enoxacin
Flucytosine
Griseofulvin
Lomefloxacin
Minocycline
Nalidixic Acid
Narfloxacin
Ofloxacin
Oxytetracycline
Pyrazinamide
Sulfonamides
Tetracycline

Antidepressants

Amitriptyline
Amoxapine
Clomipramine
Desipramine
Doxepin
Imipramine
Maprotiline
Nortriptyline
Phenelzine
Protriptyline
Trazodone
Trimipramine

Antipsychotic

Chlorpromazine
Fluphenazine
Haloperidol
Perphenazine
Prochlorperazine
Thioridazine
Thiothixane
Trifluoperazine
Trifluoperazine
Thioflupromazine

Antihypertensives

Captopril
Diltiazem
Methyldopa
Minoxidil
Nifedipine

Antiparasitic

Chloroquine
Quinine
Thiabendazole

Diuretics

Acetazolamide
Amiloride
Bendroflumethiazide
Chlorothiazide
Furosemide
Hydrochlorothiazide
Hydroflumethiazide
Methclothiazide